

University of Groningen

Costs and Effects of Abdominal versus Laparoscopic Hysterectomy

Bijen, Claudia B. M.; Vermeulen, Karin M.; Mourits, Marian J. E.; de Bock, Geertruida H.

Published in:
 PLoS ONE

DOI:
 [10.1371/journal.pone.0007340](https://doi.org/10.1371/journal.pone.0007340)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bijen, C. B. M., Vermeulen, K. M., Mourits, M. J. E., & de Bock, G. H. (2009). Costs and Effects of Abdominal versus Laparoscopic Hysterectomy: Systematic Review of Controlled Trials. *PLoS ONE*, 4(10), [7340]. <https://doi.org/10.1371/journal.pone.0007340>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Costs and Effects of Abdominal versus Laparoscopic Hysterectomy: Systematic Review of Controlled Trials

Claudia B. M. Bijen¹, Karin M. Vermeulen², Marian J. E. Mourits¹, Geertruida H. de Bock^{1*}

¹ Department of Gynecologic Oncology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, ² Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

Abstract

Objective: Comparative evaluation of costs and effects of laparoscopic hysterectomy (LH) and abdominal hysterectomy (AH).

Data sources: Controlled trials from Cochrane Central register of controlled trials, Medline, Embase and prospective trial registers.

Selection of studies: Twelve (randomized) controlled studies including the search terms costs, laparoscopy, laparotomy and hysterectomy were identified.

Methods: The type of cost analysis, perspective of cost analyses and separate cost components were assessed. The direct and indirect costs were extracted from the original studies. For the cost estimation, hospital stay and procedure costs were selected as most important cost drivers. As main outcome the major complication rate was taken.

Findings: Analysis was performed on 2226 patients, of which 1013 (45.5%) in the LH group and 1213 (54.5%) in the AH group. Five studies scored ≥ 10 points (out of 19) for methodological quality. The reported total direct costs in the LH group (\$63,997) were 6.1% higher than the AH group (\$60,114). The reported total indirect costs of the LH group (\$1,609) were half of the total indirect in the AH group (\$3,139). The estimated mean major complication rate in the LH group (14.3%) was lower than in the AH group (15.9%). The estimated total costs in the LH group were \$3,884 versus \$3,312 in the AH group. The incremental costs for reducing one patient with major complication(s) in the LH group compared to the AH group was \$35,750.

Conclusions: The shorter hospital stay in the LH group compensates for the increased procedure costs, with less morbidity. LH points in the direction of cost effectiveness, however further research is warranted with a broader costs perspective including long term effects as societal benefit, quality of life and survival.

Citation: Bijen CBM, Vermeulen KM, Mourits MJE, de Bock GH (2009) Costs and Effects of Abdominal versus Laparoscopic Hysterectomy: Systematic Review of Controlled Trials. PLoS ONE 4(10): e7340. doi:10.1371/journal.pone.0007340

Editor: Hany Abdel-Aleem, Assiut University Hospital, Egypt

Received: April 14, 2009; **Accepted:** September 9, 2009; **Published:** October 5, 2009

Copyright: © 2009 Bijen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study is funded by Zon-Mw grant (945-071-01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: g.h.de.bock@epi.umcg.nl

Introduction

Traditionally abdominal hysterectomy (AH) is the standard procedure for gynecological malignancy and several benign indications and remains the 'fallback option' if the uterus cannot be removed by another approach. While the abdominal surgical approach is an accepted effective treatment, it is associated with substantial morbidity, mostly wound problems on the short-term [1–3] and incisional hernias on the long-term [4]. A good alternative approach for patients with an indication for removal of the uterus is by laparoscopy. The laparoscopic approach to hysterectomy has evolved over the last 20 years. In 1984, Kurt Semm [5] introduced laparoscopic assistance to complicated vaginal hysterectomy and this was followed by laparoscopically assisted vaginal hysterectomy as described by Harry Reich [6]. Three types of laparoscopic hysterectomy (LH) are currently

practiced: Laparoscopically Assisted Vaginal Hysterectomy (LAVH), Total Laparoscopic Hysterectomy (TLH) and Laparoscopically Assisted Supracervical Hysterectomy (LASH). The laparoscopic approach has the advantages of laparotomy, i.e. possibility of thorough abdominal inspection to assess the abdominal cavity for extra-uterine spread and collection of peritoneal fluid for cytology. Moreover, since patients do not have a large abdominal wound, the laparoscopic approach results in a shorter hospital stay, less abdominal wound morbidity and quicker return to activity in daily life [1,2]. Nevertheless, for several reasons laparoscopy is not an established procedure for all indications for abdominal hysterectomy yet. The first reason is inexperience of surgeons with this advanced laparoscopic procedure, which results in a higher peri-operative complication rate during the learning curve [7]. Next reason is economic: higher per-operative costs, longer operation time, expensive surgical

(disposable) equipment and extra costs in converted procedures. However, laparoscopic approaches to hysterectomy offer the prospect of improved outcomes and gains in cost effectiveness through better and quicker convalescence and shorter length of inpatient stay.

A combination of the development of expensive new treatments for patients, the limited budgets available for health care and the increasing demands on the health system has led to a need to evaluate the costs in addition to clinical effects, in order to make rational decisions regarding the acceptance of new treatments into the health service.

Therefore, it is important to determine the actual costs of a new treatment and to compare those with the costs of the standard treatment, in order to give this new method a chance to be fairly judged and further propagated. With the exception of some observational studies [8–10] and small randomized trials [11,12], little is known about the costs and cost effectiveness of LH relative to the conventional abdominal approach. Our aim of this study is to pool the data of controlled trials and review whether this new health technology, laparoscopic hysterectomy, provides ‘good value for money’ in comparison to the conventional procedure.

Methods

Search strategy

The methodology of the review was according to the QUORUM statement [13] (appendix S1). Studies comparing the costs and cost effectiveness of LH versus AH were sought from a systematic review of the literature. The electronic databases Medline, Embase and the Cochrane library database were searched for relevant articles between the years 1990 and 2008. Prospective clinical trial registers were also searched for the same keywords. Search terms used were: costs, laparoscopy, laparotomy and hysterectomy. The references of all relevant articles were hand-searched for any previously missed articles. The results of the search in electronic databases and the hand search were indicated in the flowchart (figure 1).

Selection of trials

All studies that compared costs between AH and all forms of LH were included in this review. Studies were excluded from the review if they made comparisons other than those specified above. There was no restriction to any language. Furthermore, reviews, letters, author’s replies, retrospective studies and case series with no control group were excluded, whereas prospective controlled studies and randomized controlled trials were included. Papers were independently appraised by two authors (KV, CB) in terms of their methodology and design by using a predetermined protocol and graded as to the level of evidence (table 1 and 2). Differences were resolved by discussion between the authors.

The search strategy identified 16 studies [11,12,14–27] that compared the costs of laparoscopic hysterectomy and abdominal hysterectomy. Four [23–25,27] of these were not suitable for further reviewing according to our in- and exclusion criteria. The flow of studies included in the analysis is presented in figure 1 (based on the QUORUM flowchart) [13].

Data management

All data were extracted independently by two reviewers (KV, CB) and differences of opinion were resolved by consensus. Extracted data are presented in detail in appendix S2 and general study characteristics are shown in table 2. The economic impact of a surgical procedure can be divided into direct medical costs and indirect costs. Direct costs relate to health care costs and include

those relating to the hospital stay, operative procedure and treatment-related complications. Procedure costs relate to the instruments and equipment used for the procedure, the duration (including anesthetic time), staff costs and overheads costs of the theatre. Indirect costs are the societal (because of a loss of productivity) and individual costs relating to a patient’s absence from work or normal activities; this also includes costs of carers. Wherever possible, costs have been classified in this manner, although some of the studies vary in their definitions.

Analysis of (inter study) costs and effects

Reported costs. The type of cost analysis, perspective of cost analyses and all separate cost components of the studies were assessed (table 2 and 3). In case a vaginal hysterectomy arm was included, these data were excluded from further analysis. The direct and indirect costs as reported by the authors between the procedures and within studies were specified and expressed in US dollar (table 3). No formal meta-analysis could be performed because of the heterogeneity in the selected studies in terms of date of publication, currency in which costs were calculated, statistical method used, disease process examined and particularly the range of cost components that were included and primary effect measures which were taken into account.

Estimated costs and effects. To be able to draw a global conclusion with regard to costs and the ratio between costs and effects, overall treatment costs were estimated based on costs in the current situation in Dutch hospitals (table 4 and 5). Costs of the procedure were estimated based on personnel costs (\$4.14 or €3.12/min), overhead and housing (45%) according to the Dutch guidelines for costs studies. Costs of disposables (\$1,328 or €1,000: only LH) were globally assessed based on the current Dutch protocol [28]. Costs of hospital stay were based on Dutch standard prices (mean of general hospital and university medical center: \$571/€430). The price level used was that of 2007 and costs are expressed in US dollars (\$). Since information on other cost categories was scarce in most studies, those costs were not included in the present estimation. As primary effect measure the overall complication rate as well as the major complication rate was taken. The division between and definition of major and minor complications were subtracted from the original article. The complication rate consists of the reported complications added up with the amount of re-interventions performed. Conversions to laparotomy were not considered as complications.

Statistical analysis. All the analyses were done for all included studies together and a sub-analysis is done for the RCTs only. Total costs were calculated for each treatment arm in each study separately. Mean estimated costs with the range (minimum–maximum) were presented. To compare cost and effect pairs between studies, both costs and results were re-calculated to the level of 100 patients per treatment arm. Subsequently, point estimates of cost and effect pairs were plotted in a cost-effectiveness plane. These analyses were performed in Microsoft Office Excel (2003).

Results

Characteristics of selected studies

Of all included trials, 6 of the 12 controlled trials concerned randomized controlled trials [11,12,17,20–22]. Five studies scored ≥ 10 points (out of 19) for methodological quality. In the majority of studies (7 out of 12) the costs were analyzed with a cost consequence analysis, mostly from a hospital perspective. No influence of funding or sponsoring was reported. In only two studies, (also) women with a malignant indication for a




Figure 1. Flowchart of included studies.

doi:10.1371/journal.pone.0007340.g001

hysterectomy were recruited. The interventions performed in the trials varied from a simple hysterectomy to a more extended hysterectomy with or without lymphadenectomy. Furthermore, the extent to which the hysterectomy was completed laparoscopically differed from a LAVH to a LSH or a TLH. In total, 2902 patients were included, of which 662 patients underwent a vaginal hysterectomy or were allocated to a vaginal arm. Analysis was performed on 2226 patients; 1213 (54.5%) in the LH group and 1013 (45.5%) in the AH group.

The total direct costs of the LH group (\$63,997) were 6.1% higher than the AH group (\$60,114). Three studies reported lower costs for the LH group compared with the AH group [15,18,20]. The total indirect costs of the LH group (\$1,609) were half of the

total indirect costs in the AH group (\$3,139). A direct relation between costs and included cost components could not be observed.

Cost components: procedure and hospital stay

From table 4 can be concluded that the mean duration of a laparoscopic procedure ranges from 81.9 minutes [12] to 200.6 minutes [15]. The weighted mean duration of a laparoscopic procedure was 127.7 minutes. In two studies other procedures were performed in addition to the hysterectomy, equally distributed over both arms [18,19]. Conversely, the shortest mean duration of an AH was 47.3 minutes [12] and the longest mean duration 146.0 minutes [22] (table 4). The weighted mean duration of an AH was 101.3 minutes. The shortest mean hospital

Table 1. Methodological quality assessment*.

Category	Subcategory	score
<i>Generation of allocation sequence</i>	Computer generated random number	2
	Not described	1
<i>Allocation concealment</i>	Central randomization	3
	Sealed enveloped or similar	2
	Not described or inadequate	1
<i>Outcome assessor blindness</i>	Blinded for treatment arm	2
	Inadequate blinding	1
	Not described or no double-blinding	0
<i>Description of withdrawals and dropouts</i>	Numbers and reasons are described	2
	Number or reasons are not described	1
	Numbers and reasons are not described	0
<i>Efficacy of randomization/controlling</i>	Pretreatment variables in tabular form	2
	Balance of pretreatment variables mentioned but not in tabular form	1
	No information report	0
<i>Analysis include an intention-to treat analysis</i>	Intention-to treat analysis described	2
	Unclear	1
	No intention-to-treat analysis used	0
<i>All important and relevant costs for each arm</i>	For direct and indirect costs	2
	For direct or indirect costs	1
	Incomplete cost analysis	0
<i>Valid measurements and detailed description of costs</i>	Yes	2
	Unclear	1
	No	0
<i>Relevant sensitivity analysis</i>	Sensitivity analysis described and in tabular/graphical form	2
	Sensitivity analysis described, no tabular/graphical form	1
	No sensitivity analysis described	0

* = analogously of Jadad score

doi:10.1371/journal.pone.0007340.t001

stay was 1.5 days after a LH and 2.5 days after an AH. The longest mean hospital stay was 5.1 days and 7.4 days after a LH or an AH, respectively. The weighted mean hospital stay after a LH was 3.6 days (table 4), whereas the weighted mean hospital stay after an AH was 4.8 days (table 5). Overall, in each study the mean hospital stay was lower in the LH group compared to the AH group.

Effect components: overall and major rate

Overall, the mean complication rate was 15.3% in the LH group and 21.7% in the AH group, with a complication rate difference of 6.4% (table 4 and 5). The mean major complication rate was lower in the LH group (14.3%) than in the AH group (15.9%), with a complication rate difference of 1.6%. An even stronger trend can be seen from the comparison of minor complication, 11.0% in the LH group and 23.0% in the AH, respectively. For the sub-analysis of only RCTs the mean complication rate was 19.9% in the LH group and 24.6% in the AH group, with a complication rate difference of 4.7%. The mean major complication rate was even higher in the LH group (17.8%) compared to the AH group (16.1%), with a complication difference of -1.7% (table 4 and 5).

Incremental costs and effects

The estimated mean costs for the procedure were higher in the LH group \$2,226 (\$1,807–\$2,531) than in the AH group

\$648 (\$270–\$909). In contrast, mean costs for hospital stay were lower in the LH group (\$1,658: \$856–\$2,795) compared to the AH group (\$2,664: \$1,426–\$3,424). In sum, the total costs in the LH group were \$3,884 (\$3,130–\$4,928) and in the AH group \$3,312 (\$2,208–\$3,767), with a cost difference of \$572 between groups. The incremental costs for reducing one patient with complication(s) in the LH group compared to the AH group were \$8,938, ($\$572 \times 100 / 6.4$) re-calculated to a level of 100 patients. When subdivided for major complications, the incremental costs were \$35,750 ($\$572 \times 100 / 1.6$) for reducing one patient with major complication(s). For the sub-analysis of only RCTs, the total costs in the LH group were \$3,794 (\$3,263–\$4,451) and in the AH group \$3,277 (\$2,208–\$3,767), with a cost difference of \$517. The incremental costs for reducing one patient with complication(s) in the LH group compared to the abdominal group were \$11,000 ($\$517 \times 100 / 4.7$). Regarding major complications, effects were lower with higher costs in the LH group. Three point estimates are located in the north-east quadrant of figure 2a, indicating that these studies generated extra effects of LH for relatively low additional costs, within a cost ratio range of \$2,895 to \$3,552 [11,18,22]. With regard to overall complications, six studies are located in the north-east quadrant [11,12,14,15,18,22] and therefore generating extra effects of LH within a cost ratio range of \$1,401 to \$6,618 (figure 2b).

Table 2. Study characteristics.

Author	Design	Cost analysis	Cost Perspective	Participants (n =)	Indication	Interventions	Quality assessment score
<i>Abdelmonem 2006</i>	CCT; Multi centre	Cost consequence	Hospital	177	Benign	AH LH [LAVH, LSH, TLH]	5
<i>Van den Eeden 1998</i>	CCT; Multi centre	Cost consequence	Hospital	287	Benign/Malignant	VH AH LH [LAVH]	9
<i>Ellstrom 1998</i>	RCT; Single centre	Cost consequence	Societal	143	Benign	AH +/- BSO LH [TLH] +/- BSO	13
<i>Eltabbakh 2001</i>	CCT; Single centre	Cost consequence	Hospital	147	Malignant	AH +BSO +/- LN LH [LAVH] +BSO +/- LN	10
<i>Falcone 1999</i>	RCT; Single centre	Cost minimization	Hospital	48	Benign	AH LH [LAVH]	11
<i>Howard 1993</i>	CCT; Single centre	Cost consequence	Hospital	30	Benign	AH LH [LAVH]	2
<i>Kung 1996</i>	CCT; Single centre	Cost	Hospital/Insurance	301	Benign	AH LH [LAVH]	6
<i>Leng 2004</i>	CCT; Single centre	Cost	Societal	54	Benign	AH VH	5
<i>Lumsden 2000</i>	RCT; Multi centre	Cost consequence	Hospital	190	Benign	LH AH LH [LAVH]	17
<i>Raju 1994</i>	RCT; Single centre	Cost minimization	Hospital	80	Benign	AH +BSO LH [LAVH] +BSO	7
<i>Sculpher 2004</i>	RCT; Multi centre	Cost utility	Health care provider	1380	Benign	VH AH LH [LAVH, LSH, TLH]	15
<i>Summit 1998</i>	RCT; Multi centre	Cost	Hospital	65	Benign	AH LH [LAVH]	11

CCT = controlled clinical trial; RCT = randomized controlled trial; AH = abdominal hysterectomy; LH = laparoscopic hysterectomy; LAVH = laparoscopic assisted vaginal hysterectomy; LSH = laparoscopic supracervical hysterectomy; TLH = total laparoscopic hysterectomy; VH = vaginal hysterectomy; BSO = bilateral salpingo-oophorectomy; LN = lymph nodes.
doi:10.1371/journal.pone.0007340.t002

Table 3. Actual direct and/or indirect costs per treatment arm.

Author	Currency	LH		AH		Cost components
		direct	indirect	direct	indirect	
<i>Abdelmonem 2006</i>	US dollar	\$16,451 ↑	n.i.	\$15,145	n.i.	Hospital cost, anesthesia
<i>V. d. Eeden 1998</i>	US dollar	\$8,099 ↓	n.i.	\$9,135	n.i.	Preoperative costs, hospital costs (procedure, hospital stay, physician fee), postoperative costs.
<i>Ellstrom 1998</i>	SEK	\$3,169 ↑	\$1,411 ↓	\$3,116	\$2,838	Surgery, services from other departments, hospital stay, postoperative visits, sick leave
<i>Elkabbakh 2001</i>	US dollar	\$13,051 ↑	n.i.	\$11,028	n.i.	Surgeon's fee, anesthesiologist's fee, operating room cost, hospital cost
<i>Falcone 1999</i>	US dollar	\$-277 ↑ *	n.i.	\$0 *	n.i.	Operating room cost, anesthesia, postoperative cost
<i>Howard 1993</i>	US dollar	\$3,926 ↓	n.i.	\$4,524	n.i.	Operating equipment, operating time, anesthesia, non operating costs
<i>Kung 1996</i>	NT dollar	\$1,819 ↑	n.i.	\$1,566	n.i.	Laboratory test, examination charges, surgery and material fees, anesthesia, nursing care, ward fee
<i>Leng 2004</i>	Yuan	\$876 ↑	\$198 ↓	\$749	\$301	Operating costs, outpatient department visits, readmissions, hospital stay (incl. medication), sick leave, transport costs
<i>Lumsden 2000</i>	UK pound	\$3,462 ↑	n.i.	\$2,733	n.i.	Operation costs, hospital stay, readmissions
<i>Raju 1994</i>	UK pound	\$1,909 ↓	n.i.	\$2,652	n.i.	Disposables, hospital stay, operating time
<i>Sculpher 2004</i>	UK pound	\$2,797 ↑	n.i.	\$2,492	n.i.	Ward fee, theatre (staff, overhead, disposables), visits (GP, outpatient department)
<i>Summit 1998</i>	US dollar	\$8,161 ↑	n.i.	\$6,974	n.i.	Not specified
Subtotal RCT		\$19,775 ↑	\$1,411 ↓	\$17,967	\$2,838	
Total		\$63,997 ↑	\$1,609 ↓	\$60,114	\$3,139	

n.i. = not included; * = difference in means; Exchange rates of year of performing study (<http://www.x-rates.com/cgi-bin/hlookup.cgi>); doi:10.1371/journal.pone.0007340.t003

Table 4. Costs and effects of laparoscopic hysterectomy.

Author	Costs		Effects		
	Procedure (time in minutes)	Hospital stay (days)	Complication rate * (Major)	Complication rate * (Minor)	Complication rate * (Overall)
Abdelmonem 2006	157.8	1.5	9.8 (5/51)	7.8 (4/51) ^a	13.7 (7/51)
Van den Eeden 1998	200.6	1.8	0.0 (0/56)	0.0 (0/56)	0.0 (0/56) [‡]
Ellstrom 1998	148.0	2.5	26.8 (19/71)**	n.s.	26.8 (19/71)
Elabbakh 2001	190.5	2.5	11.6 (10/86)	n.s.	11.6 (10/86)
Falcone 1999	180.0	1.5	39.1 (9/23)	n.s.	39.1 (9/23)
Howard 1993	169.0	3.7	13.3 (2/15)	n.s.	13.3 (2/15)
Kung 1996	134.0	4.9	15.2 (21/138)	n.s.	15.2 (21/138)
Leng 2004	n.a.	5.1	n.a.	n.a.	n.a.
Lumsden 2000	81.9	4.0	5.3 (5/95)	n.s.	8.0
Raju 1994	100.0	3.5	5.0 (2/40)	n.s.	5.0 (2/40)
Sculpher 2004	108.1	4.0	10.1 (59/584)**	25.2 (147/584)**	n.s.
Summit 1998	179.8	2.1	20.6 (7/34)	n.s.	20.6 (7/34) [‡]
Weighted mean RCT	113.0	3.7	17.8	25.2	19.9
Weighted mean	127.7	3.6	14.3	11.0	15.3

n.s. = not specified; n.a. = not available; * = percentage of patients with complications; ** = data derived from original article; ^a = converted patients reported on complication level instead of patient level. Therefore these patients can not be excluded; [‡] = number of complications reported.
doi:10.1371/journal.pone.0007340.t004

Table 5. Costs and effects of abdominal hysterectomy.

Author	Costs		Effects		
	Procedure (time in minutes)	Hospital stay (days)	Complication rate * (Major)	Complication rate * (Minor)	Complication rate * (Overall)
Abdelmonem 2006	127.2	3.7	10.0 (5/50)	28.0 (14/50)	32.0 (16/50)
Van den Eeden 1998	151.4	3.3	1.2 (2/164)	14.0 (23/164)	15.2 (25/164)‡
Ellstrom 1998	93.1	5.0	33.3 (24/72)**	n.s.	33.3 (24/72)**
Elkabbakh 2001	132.8	5.2	8.8 (5/57)	n.s.	8.8 (5/57)
Falcone 1999	130.0	2.5	23.8 (5/21)	n.s.	23.8 (5/21)
Howard 1993	119.0	5.2	40.0 (6/15)	n.s.	40.0 (6/15)
Kung 1996	112.0	5.2	17.8 (28/157)	n.s.	17.8 (28/157)
Leng 2004	n.a.	7.4	n.a.	n.a.	n.a.
Lumsden 2000	47.3	5.7	1.1 (1/95)	n.s.	14.0
Raju 1994	57.0	6.0	0.0 (0/40)	n.s.	0.0 (0/40)
Sculpher 2004	74.1	5.1	6.2 (18/292)*	27.1 (79/292)**	n.s.
Summit 1998	146.0	4.1	32.3 (10/31)	n.s.	32.3 (10/31)‡
Weighted mean RCT	76.9	5.1	16.1	27.1	24.6
Weighted mean	101.3	4.8	15.9	23.0	21.7

n.s. = not specified; n.a. = not available; * = percentage of patients with complications; ** = data derived from original article; ‡ = number of complications reported.
doi:10.1371/journal.pone.0007340.t005





Figure 2. Incremental costs and effects for patients with major complications (A) and all complications (B) per 100 patients ■ = RCT □ = CCT.
doi:10.1371/journal.pone.0007340.g002

Discussion

For the first time, costs and short term effects were reviewed and calculated between laparoscopic hysterectomy and abdominal hysterectomy in a large group of patients. Cost evaluations for other diseases comparing laparoscopic and open surgery already published, demonstrated that laparoscopic appendectomy, laparoscopic colectomy and laparoscopic cholecystectomy results in decreased hospital stay, lower hospital cost and faster return to work or daily activities than open procedures [29]. Our review showed a similar trend with regard to hospital stay and hospital costs. Twelve trials concerning 2226 patients in total were reviewed, of which five trials scored ≥ 10 points (out of 19) for methodological quality. As reported by the authors, the direct costs were higher and indirect costs were lower for LH compared to AH. Procedure costs were estimated higher and costs of hospital stay were estimated lower in favor of LH. In the LH group the estimated overall and major complication rate were both lower compared to the AH group, 6.4% and 1.6% respectively. Overall, the incremental cost for reducing one patient with complication(s) in the LH group compared to the AH group was \$8,938 and \$35,750 for reducing one patient with major complication(s).

Based on this data, it is not clear which costs are really associated with the actual cost difference between both treatment modalities. There is significant variability between studies in the range of cost components covered. Although some authors give reasonable detail on economic data, [11,12,15,21] many give scanty information or do not even specify the method of measuring costs [14,16–20,22]. Strikingly, total costs were far lower for the Chinese and Taiwanese study [19,26] compared to the other included studies. Presumably, costs for hospital stay and/or out patient department visits were much lower in Asia than in Western countries. The direct costs were higher for LH, whereas the indirect costs were lower than an AH. Nevertheless, only one study collected data on indirect costs. Importantly, previous reports stated that there might be societal benefits associated with lower indirect cost for LH [7,16,22,30–34]. In this analysis, cost drivers considered were the procedure (including use of disposables) and hospital stay. Noticeably, there is a wide range in these main cost drivers between the studies. The wide range is caused by the duration of operation time and the experience of the surgeons. Operating time is related to the disease process (e.g. benign or malignant) and the occurrence of additional procedures. A more extended hysterectomy in case of a malignancy might have influence on the length of hospital stay, as well. Inevitably, some of the studies have compared inexperienced laparoscopic surgeons with experienced ‘open’ surgeons. As surgeons become more experienced with laparoscopic procedures, the length of operating time decreases [35–37]. By training of surgeons in laparoscopy and using reusables instead of expensive disposables, procedure costs for LH can be drastically reduced.

The same diversity as is found in costs between studies can be seen in complication rates. In two studies [15,22] an overestimation of complication rate might be shown in table 4, since only number of complications were reported and not number of patients with complications. Some studies reported major and minor complications separately [12,14,15,21], with high minor complication rates. A possible explanation for this phenomenon can be the definition of minor or major complications. Standardized criteria for defining minor and major complications should be used in order to adequately compare complication rate [38]. In most of the studies postoperative complications and short-

term post operative complications were reported and not long-term complications. We therefore could not specify the occurrence of long-term complications. Besides, the studies were selected on bases of including costs in their report, not on morbidity as effect measure. Since a part of the selected studies were of low methodological quality (< 10 points), the reported morbidity rates might dissent from well-designed studies, powered on morbidity. In comparison, in a large randomized controlled trial Garry et al. demonstrated a major complication rate of 9.6% for LH and 6.2% for AH, excluding conversions regarded as complications [39]. Reasons to convert a procedure are to avoid a complication to occur or because of a complication that occurred. In the latter, the complication is already recorded separately. Calculating a mean effect, while some studies showed an opposite effect might give a distorted picture. Again, reasons for this diversity are the experience of the surgeons and the indication of performing a hysterectomy. Inexperience with this advanced laparoscopic procedure results in a higher peri-operative complication rate during the learning curve [7].

The sub analyses of only RCTs indicated that the amount of costs were higher or did not show additional effects when looking at major complication rate. The latter means that implementation of the laparoscopic technique is not cost effective as a standard procedure. Minimization of selection bias and information bias might be explanations for the higher morbidity rates in the LH group and as a result not being cost effective as was found in RCTs compared to all controlled trials. Randomization is important as it helps reduce the possibility of bias [40]. Without randomization, there may be a tendency for researchers to select these participants for particular intervention groups they favor (i.e. selection bias). In this way, systematically different estimates of treatment effects can be yielded [41]. Moreover, most of the RCTs were rigorously set up and monitored to ensure accurate registration of complications and therefore safeguarding a systematic bias in reporting frequency (i.e. information bias) by approach.

However, based on other criteria such as quality of life and survival, LH might be more advantageous than AH. Results of two large RCTs i.e. the GOG Lap2 study [42] and the LACE trial [43] are expected in the near future with quality of life and survival as primary endpoints. This will make more definitive conclusions about these specific outcomes possible. Since the aim of the present review was to globally explore differences in costs and effects between both treatment arms independent of the indication, the analyses of all controlled trials were more important and decisive. Furthermore, because the relative differences were analyzed instead of absolute differences the indication for removal of the uterus and consequently the extent of the procedure was therefore considered less relevant.

On average, an amount of \$8,938 has to be invested to reduce one complication when performing a LH in stead of an AH. The explicit quantification of acceptable costs for a given benefit is difficult to define. In other words, what costs are acceptable for reducing one complication? From previous cost reports, a cut of value of \$100,000 for reducing one death seems cost effective and therefore acceptable [44]. No reports are available, defining a cut off value for reduction one complication. The National Institute for Health and Clinical Excellence (NICE) has adopted a cost effectiveness threshold range of \$40,000–\$60,000 per Quality Adjusted Life Year (QALY) [45]. According to this guideline a conservative range of \$0–\$20,000 to prevent one additional complication represents an acceptable cost effectiveness ratio, considering the (unknown) variation of complications and the consequences of a complication in terms of prolonged hospital stay, re-intervention and patient burden.

Conclusion

The perception that laparoscopic procedures are more costly than open procedures has been a major reason for the slow acceptance of laparoscopic surgery. Important is whether LH is more effective and more costly, with the added benefit worth the added costs. From this study it can be concluded that the benefits of shorter hospital stay in the LH group might compensate for the increased procedure costs. Laparoscopic hysterectomy points in the direction of cost effectiveness, however a broader perspective is needed, including indirect costs and long term effects as survival and quality of life for determining implications for practice. Currently, a cost effectiveness evaluation of the two surgical approaches is being conducted alongside a large multi-centre RCT [46], complying with all guidelines.

Supporting Information

Appendix S1 QUORUM checklist

References

- Manolitsas TP, McCartney AJ (2002) Total laparoscopic hysterectomy in the management of endometrial carcinoma. *J Am Assoc Gynecol Laparosc* 9: 54–62.
- Obermair A, Manolitsas TP, Leung Y, Hammond IG, McCartney AJ (2005) Total laparoscopic hysterectomy versus total abdominal hysterectomy for obese women with endometrial cancer. *Int J Gynecol Cancer* 15: 319–324.
- Gemignani ML, Curtin JP, Zelmanovich J, Patel DA, Venkatraman E, et al. (1999) Laparoscopic-assisted vaginal hysterectomy for endometrial cancer: clinical outcomes and hospital charges. *Gynecol Oncol* 73: 5–11.
- Franchi M, Ghezzi F, Buttarelli M, Tateo S, Balestreri D, et al. (2001) Incisional hernia in gynecologic oncology patients: a 10-year study. *Obstet Gynecol* 97: 696–700.
- Semm K (1984) [Endoscopic methods in gastroenterology and gynecology. Completion of diagnostic pelviscopy by endoscopic abdominal surgery]. *Fortschr Med* 102: 534–537.
- Reich H (1989) New techniques in advanced laparoscopic surgery. *Baillieres Clin Obstet Gynaecol* 3: 655–681.
- Garry R, Fountain J, Brown J, Manca A, Mason S, et al. (2004) EVALUATE hysterectomy trial: a multicentre randomised trial comparing abdominal, vaginal and laparoscopic methods of hysterectomy. *Health Technol Assess* 8: 1–154.
- Lowell L, Kessler AA (2000) Laparoscopically assisted vaginal hysterectomy. A suitable substitute for abdominal hysterectomy? *J Reprod Med* 45: 738–742.
- Schneider A, Merker A, Martin C, Michels W, Krause N (1997) Laparoscopically assisted vaginal hysterectomy as an alternative to abdominal hysterectomy in patients with fibroids. *Arch Gynecol Obstet* 259: 79–85.
- Chapron C, Fernandez B, Dubuisson JB (2000) Total hysterectomy for benign pathologies: direct costs comparison between laparoscopic and abdominal hysterectomy. *Eur J Obstet Gynecol Reprod Biol* 89: 141–147.
- Ellstrom M, Ferraz-Nunes J, Hahlin M, Olsson JH (1998) A randomized trial with a cost-consequence analysis after laparoscopic and abdominal hysterectomy. *Obstet Gynecol* 91: 30–34.
- Lumsden MA, Twaddle S, Hawthorn R, Traynor I, Gilmore D, et al. (2000) A randomised comparison and economic evaluation of laparoscopic-assisted hysterectomy and abdominal hysterectomy. *BJOG* 107: 1386–1391.
- Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, et al. (1999) Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. *Quality of Reporting of Meta-analyses*. *Lancet* 354: 1896–1900.
- Abdelmonem A, Wilson H, Pasic R (2006) Observational comparison of abdominal, vaginal and laparoscopic hysterectomy as performed at a university teaching hospital. *J Reprod Med* 51: 945–954.
- Van Den Eeden SK, Glasser M, Mathias SD, Colwell HH, Pasta DJ, et al. (1998) Quality of life, health care utilization, and costs among women undergoing hysterectomy in a managed-care setting. *Am J Obstet Gynecol* 178: 91–100.
- Eltabbakh GH, Shamonki MI, Moody JM, Garafano LL (2001) Laparoscopy as the primary modality for the treatment of women with endometrial carcinoma. *Cancer* 91: 378–387.
- Falcone T, Paraiso MF, Mascha E (1999) Prospective randomized clinical trial of laparoscopically assisted vaginal hysterectomy versus total abdominal hysterectomy. *Am J Obstet Gynecol* 180: 955–962.
- Howard FM, Sanchez R (1993) A comparison of laparoscopically assisted vaginal hysterectomy and abdominal hysterectomy. *J Gynecol Surg* 9: 83–90.
- Kung FT, Hwang FR, Lin H, Tai MC, Hsieh CH, et al. (1996) Comparison of laparoscopically assisted vaginal hysterectomy and abdominal hysterectomy in taiwan. *J Formos Med Assoc* 95: 769–775.
- Raju KS, Auld BJ (1994) A randomised prospective study of laparoscopic vaginal hysterectomy versus abdominal hysterectomy each with bilateral salpingo-oophorectomy. *Br J Obstet Gynaecol* 101: 1068–1071.
- Sculpher M, Manca A, Abbott J, Fountain J, Mason S, et al. (2004) Cost effectiveness analysis of laparoscopic hysterectomy compared with standard hysterectomy: results from a randomised trial. *BMJ* 328: 134.
- Summitt RL, Jr., Stovall TG, Steege JF, Lipscomb GH (1998) A multicenter randomized comparison of laparoscopically assisted vaginal hysterectomy and abdominal hysterectomy in abdominal hysterectomy candidates. *Obstet Gynecol* 92: 321–326.
- Bornstein SJ, Shaber RE (1995) Laparoscopically assisted vaginal hysterectomy at a health maintenance organization. Cost-effectiveness and comparison with total abdominal hysterectomy. *J Reprod Med* 40: 435–438.
- Doucette RC, Scott JR (1996) Comparison of laparoscopically assisted vaginal hysterectomy with abdominal and vaginal hysterectomy. *J Reprod Med* 41: 1–6.
- Kunz G, Plath T, Leyendecker G (1996) [Comparison of laparoscopically assisted vaginal hysterectomy with abdominal hysterectomy. Technique and results]. *Geburtshilfe Frauenheilkd* 56: 453–457.
- Leng JH, Lang JH, Li HJ, Liu ZF, Sun DW, et al. (2004) [Effect of different surgical approaches on quality of life and cost-consequence]. *Zhonghua Fu Chan Ke Za Zhi* 39: 315–318.
- Raju KS, Auld BJ (1994) Laparoscopic-assisted vaginal hysterectomy with bilateral oophorectomy versus total abdominal hysterectomy and bilateral salpingo-oophorectomy—implications for gynecologic practice in the United Kingdom. *J Am Assoc Gynecol Laparosc* 1: 111–115.
- Oostenbrink JB, Bouwmans CAM, Koopmanschap MA, Rutten FFH (2004) Guidelines for cost-studies: methods and unit prices for economic evaluations in health care, revised version Dutch Health insurance board.
- Roumm AR, Pizzi L, Goldfarb NI, Cohn H (2005) Minimally invasive: minimally reimbursed? An examination of six laparoscopic surgical procedures. *Surg Innov* 12: 261–287.
- Holub Z, Bartos P, Dorr A, Eim J, Jabor A, et al. (1999) The role of laparoscopic hysterectomy and lymph node dissection in treatment of endometrial cancer. *Eur J Gynaecol Oncol* 20: 268–271.
- Harkki-Siren P, Sjöberg J, Toivonen J, Tiitinen A (2000) Clinical outcome and tissue trauma after laparoscopic and abdominal hysterectomy: a randomized controlled study 1. *Acta Obstet Gynecol Scand* 79: 866–871.
- Olsson JH, Ellstrom M, Hahlin M (1996) Cost Effectiveness and Health Status after Laparoscopic and Abdominal Hysterectomy. *J Am Assoc Gynecol Laparosc* 3: S37.
- Seracchioli R, Venturoli S, Vianello F, Govoni F, Cantarelli M, et al. (2002) Total laparoscopic hysterectomy compared with abdominal hysterectomy in the presence of a large uterus. *J Am Assoc Gynecol Laparosc* 9: 333–338.
- Langebrekke A, Eraker R, Nesheim BI, Urnes A, Busund B, et al. (1996) Abdominal hysterectomy should not be considered as a primary method for uterine removal. A prospective randomised study of 100 patients referred to hysterectomy. *Acta Obstet Gynecol Scand* 75: 404–407.
- Holub Z, Jabor A, Bartos P, Hendl J, Urbanek S (2003) Laparoscopic surgery in women with endometrial cancer: the learning curve. *Eur J Obstet Gynecol Reprod Biol* 107: 195–200.
- Eltabbakh GH (2000) Effect of surgeon's experience on the surgical outcome of laparoscopic surgery for women with endometrial cancer. *Gynecol Oncol* 78: 58–61.
- Perino A, Cucinella G, Venezia R, Castelli A, Cittadini E (1999) Total laparoscopic hysterectomy versus total abdominal hysterectomy: an assessment of the learning curve in a prospective randomized study. *Hum Reprod* 14: 2996–2999.

Found at: doi:10.1371/journal.pone.0007340.s001 (0.03 MB PDF)

Appendix S2 Characteristics of included studies

Found at: doi:10.1371/journal.pone.0007340.s002 (0.15 MB DOC)

Acknowledgments

We thank Ms. W. Lu for her assistance with translating the Chinese manuscript and Ms. N. Leffers for constructing the cost effect planes.

Author Contributions

Analyzed the data: CB KV GdB. Wrote the paper: CB. Contributed to the study design and data interpretation: CB KV MM GdB. Checked the first draft for important intellectual content and provided many valuable comments: KV MM GdB. Approved the final version of the manuscript: CB KV MM GdB. Data collection was undertaken by: CB. Contributed to the selection and extraction of data from the included studies: CB KV.

38. National Cancer Institute (2009) CTEP, NCI guidelines: Adverse events reporting requirements. .
39. Garry R, Fountain J, Mason S, Hawe J, Napp V, et al. (2004) The eVALuate study: two parallel randomised trials, one comparing laparoscopic with abdominal hysterectomy, the other comparing laparoscopic with vaginal hysterectomy 1. *BMJ* 328: 129.
40. Sibbald B, Roland M (1998) Understanding controlled trials. Why are randomised controlled trials important? *BMJ* 316: 201.
41. Schulz KF, Chalmers I, Hayes RJ, Altman DG (1995) Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 273: 408–412.
42. GOG-LAP2 (2009) Gynecologic Oncology Group National Cancer Institute (NCI). Laparoscopic surgery or standard surgery in treating patients with endometrial cancer or cancer of the uterus. CDR0000064513, GOG-LAP2.
43. Janda M, Gebiski V, Forder P, Jackson D, Williams G, et al. (2006) Total laparoscopic versus open surgery for stage 1 endometrial cancer: the LACE randomized controlled trial. *Contemp Clin Trials* 27: 353–363.
44. Subak LL, Caughey AB (2000) Measuring cost-effectiveness of surgical procedures. *Clin Obstet Gynecol* 43: 551–560.
45. Appleby J, Devlin N, Parkin D (2007) NICE's cost effectiveness threshold 2. *BMJ* 335: 358–359.
46. Bijen CB, Briet JM, de Bock GH, Arts HJ, Bergsma-Kadijk JA, et al. (2009) Total laparoscopic hysterectomy versus abdominal hysterectomy in the treatment of patients with early stage endometrial cancer: a randomized multi center study 2. *BMC Cancer* 9: 23.